

REVIEW

Serotonin, serotonin receptors and their actions in insects

Rut Vleugels*, Heleen Verlinden*, Jozef Vanden Broeck

Department of Animal Physiology and Neurobiology, Zoological Institute, KU Leuven, Naamsestraat 59, 3000 Leuven, Belgium

Correspondence: Heleen Verlinden

E-mail: heleen.verlinden@bio.kuleuven.be

Received: September 02, 2014

Published online: January 08, 2015

* These authors contributed equally to this paper

Serotonin is an ancient monoamine neurotransmitter, biochemically derived from tryptophan. It is most abundant in the gastrointestinal tract, but is also present throughout the rest of the body of animals and can even be found in plants and fungi. Serotonin is especially famous for its contributions to feelings of well-being and happiness. More specifically it is involved in learning and memory processes and is hence crucial for certain behaviors throughout the animal kingdom. This brief review will focus on the metabolism, biological role and mode-of-action of serotonin in insects. First, some general aspects of biosynthesis and break-down of serotonin in insects will be discussed, followed by an overview of the functions of serotonin, serotonin receptors and their pharmacology. Throughout this review comparisons are made with the vertebrate serotonergic system. Last but not least, possible applications of pharmacological adjustments of serotonin signaling in insects are discussed.

Keywords: behavior; biogenic amine; drug; G protein-coupled receptor; insecticide; learning and memory

To cite this article: Rut Vleugels, et al. Serotonin, serotonin receptors and their actions in insects. Neurotransmitter 2015; 2: e314. doi: 10.14800/nt.314.

Introduction

Serotonin (5-hydroxytryptamine or 5-HT) is an ancient monoamine neurotransmitter that belongs to the biogenic amines. Biogenic amines are derived from aromatic amino acids and function as neurotransmitters and neuromodulators in the central nervous system or can be released in circulation to act as (neuro)hormones. In vertebrates there are five established biogenic amine neuromodulators: the indolamine 5-HT, the imidazolamine histamine and the catecholamines dopamine, epinephrine (also called adrenaline) and norepinephrine (also called noradrenaline). These biogenic amines are often grouped with other small compounds, such as acetylcholine, glutamate and gamma-aminobutyric acid, as small molecule neurotransmitters. The phenolamines tyramine and octopamine are generally considered as the protostome analogs of the adrenergic amines in vertebrates and other deuterostomes^[1-4]. However, octopaminergic systems have also been reported in some deuterostomian phyla (in species, such as the echinoderm sunflower star

Psyconopodia helanthoides and the cephalochordate *Branchiostoma lanceolatum*) and monophenolic amines are likely to occur in considerable quantities in all triblastic animals, except vertebrates^[2,5,6]. In vertebrates, octopamine and tyramine are present in only very small quantities (trace amines)^[7].

Biosynthesis of 5-HT

Biogenic amines are synthesized starting from amino acids by only one or a few enzymatic steps, including a decarboxylation. 5-HT belongs to the class of monoamines, where a basic amine group is separated from an aromatic core by an aliphatic chain with two carbon oxides. The synthesis of 5-HT begins with the essential amino acid tryptophan. The enzyme tryptophan hydroxylase (TPH), also known as tryptophan-5-monooxygenase, first adds a hydroxyl group to tryptophan forming 5-hydroxytryptophan (5-HTP)^[8] (Figure 1). The hydroxylase TPH constitutes together with phenylalanine hydroxylase (which catalyzes




Figure 1. Biosynthesis of serotonin (5-HT). Tryptophan is converted into 5-hydroxytryptamine (5-HTP) by tryptophan hydroxylase (TPH). Next, 5-HTP is converted to 5-HT by 5-HTP decarboxylase (also called aromatic amino acid decarboxylase or AADC).

the hydroxylation of the aromatic side-chain of phenylalanine to generate tyrosine) and tyrosine hydroxylase (which converts tyrosine to 3,4-dihydroxyphenylalanine or DOPA) the non-heme iron and tetrahydrobiopterin-dependent aromatic amino acid hydroxylase family. These enzymes use (6R)-L-erythro-5,6,7,8-tetrahydrobiopterin and O_2 as cofactors^[9,10].

In a second step of 5-HT biosynthesis, 5-HTP decarboxylase catalyzes the conversion of 5-HTP to 5-HT^[11] (Figure 1). It has been shown that 5-HTP decarboxylase is the same enzyme as dopamine decarboxylase, which catalyzes the decarboxylation step in dopamine biosynthesis, and is generally called aromatic amino acid decarboxylase^[12].

Metabolic pathways of 5-HT

To act in the nervous system, biogenic amines are stored at synaptic terminals for release into the chemical synapse in response to neuronal depolarization. In the synaptic cleft, the

amines bind specific receptor proteins (see § 5.), which can be present on both pre- and postsynaptic cells. Synaptic effects of the biogenic amines are terminated via re-uptake by binding to monoamine transporters^[13-15]. These transporter proteins belong to a subclass of the neurotransmitter/sodium symporter family, a group of glycoproteins characterized by their 12 transmembrane segments (TM) structure^[14,16]. The first identified insect amine transporter was the SERotonin Transporter (SERT) from the fruit fly *Drosophila melanogaster*^[17,18]. Later, insect transporters with high affinity for dopamine^[19] and tyramine/octopamine^[13,20-22] were also identified. In humans, these transporters attained much interest because psychoactive compounds both for medication and illegal drugs interact with them^[23].

After re-uptake, biogenic amines are repacked into vesicles for re-release or enzymatically inactivated. In vertebrates, the primary catabolic pathway for biogenic amines is oxidative deamination by monoamine oxidases^[14]. Little or no monoamine oxidase activity could be measured in insect nervous tissue^[24]. Enzymatic inactivation in invertebrates is considered to take place after amino-terminal tagging with specific groups, mainly N-acetylation or N-methylation. However γ -glutamylolation, sulphation, and β -alanyl conjugation are also possible metabolic routes^[25,26].

Functions of 5-HT in insects

5-HT is a intracellular signaling molecule found in all organisms with a central nervous system. The discovery of 5-HT as an important neuromodulator involved a multidisciplinary approach of several research groups. In the 1940s, during studies of constricting factors in the blood causing hypertension, a serum substance affecting vascular tone was isolated and named "serotonin"^[27,28]. A chemical structure elucidation approach (using ultraviolet absorption spectrometry and colorimetric chemical reactions) showed that serotonin was 5-hydroxytryptamine (5-HT)^[29]. In 1952, 5-HT was shown to be the same substance as enteramine, a compound deduced in the 1930's from an extract of the enterochromaffin cells in gastrointestinal mucosa, which caused contraction of the smooth muscle of the rat uterus^[30,31]. Approximately 80-90% of the human body's total 5-HT is located in the enterochromaffin cells of the gut, where it is used to regulate intestinal motility^[32,33]. A few years later, 5-HT was proven to be present in mammalian brain, the first step to its recognition as a neuromodulator^[34]. The observation that the hallucinatory drug lysergide acid diethylamide (LSD) antagonized a response produced by 5-HT further confirmed the idea that 5-HT may be important for the behaviour^[35].

Nowadays, the role of 5-HT is established in the regulation of many important human processes, such as appetite, mood and emotion, sleep, sexual activity, pain, and learning abilities. In different insect species, comparable processes were demonstrated to be affected by 5-HT. For example, in *D. melanogaster*, elevating 5-HT levels, either by treatment with 5-HTP or by overexpressing TPH, significantly increased periods of sleep^[36]. On the contrary, entrainment of the *Drosophila* circadian clock to light was inhibited by 5-HT^[37]. In several insect species, 5-HT was demonstrated to modulate appetite. In *D. melanogaster*, neuromodulatory actions of 5-HT were shown to depress feeding, while decreased neuronal 5-HT levels increased appetite^[38]. In another dipteran species, the flesh fly *Neobellieria bullata*, 5-HT injection in the hemolymph decreased feeding^[39]. In the honey bee, 5-HT inhibits feeding when injected in the brain and excites muscle contractions when injected in the gut, although general elevation of 5-HT in the bee's hemolymph did not affect food intake^[40]. Consistent with these findings, a 5-HT receptor was identified as a feeding modulation target from a small molecule drug screening in *Drosophila*^[41]. In the yellow-fever mosquito *Aedes aegypti*, 5-HT immunoreactive axons surround parts of the salivary gland of adult females^[42] and the blood-feeding success was reduced by administration of the 5-HT depleting drug α -methyl-tryptophan^[43]. Moreover, 5-HT increased fluid secretion from salivary glands in *A. aegypti*, the fly *Calliphora vicina* and the cockroach *Periplaneta americana*^[42,44,45]. In *Rhodnius prolixus*, 5-HT acts as a diuretic hormone. It is produced soon after the initiation of feeding and stimulates rapid tubule secretion^[46,47]. In *Drosophila*, where insulin-like peptides both regulate blood sugar and act as growth factors, serotonergic neurons control the adult body size by affecting insulin-like peptide secretion^[48,49]. How much food an animal gets is not only dependent on the abundance of food in the environment but also on the animal's social rank, since the stronger or more dominant animals may steal food from the weaker or less dominant ones. This may explain why 5-HT is not only engaged in the recognition of food availability but also of social rank.

5-HT also has thorough effects on some aspects of learning and memory and is thus a major player in modulating several insect behaviors. By inhibiting the serotonergic system in neurons of *D. melanogaster*, appetitive olfactory memory performance was considerably reduced^[50]. Flies with genetically or pharmacologically reduced 5-HT levels in the brain also had a strongly reduced memory formation in a behavioral test wherein flies were trained to avoid a chamber position associated with high temperature^[51]. In honey bees, both memory storage and retrieval were reduced when 5-HT was injected prior to

conditioning^[52-54]. In the desert locust *Schistocerca gregaria* 5-HT and its downstream effector molecules were shown to induce gregarious, swarming behavior^[55,56]. Fruit flies with genetically or pharmacologically elevated 5-HT levels showed higher fighting frequencies and more intense fighting than untreated flies^[57].

In the periphery, 5-HT seems to affect the performance of certain muscles. It was for example shown to have relaxing properties on the foregut of the migratory locust *Locusta migratoria*^[58], while other experiments demonstrated acceleration of the myogenic rhythm of the extensor-tibiae muscle in locusts by 5-HT^[59]. Also in migratory locusts, antidiuretic effects were reported for 5-HT, which may result from direct effects on the epithelial and muscular cells of the rectum^[60].

5-HT receptors

5-HT is one of the oldest signaling molecules in evolution, which may explain why 5-HT interacts with such a diversity of receptors belonging to the G protein-coupled receptor (GPCR) and the ligand-gated ion-channel families. The first 5-HT receptors are estimated to have appeared 700 to 800 million years ago in single-celled eukaryotes, such as paramecia^[61-64]. They may indeed have been amongst the first rhodopsin-like receptors reacting to a chemical^[65]. Today, the wealth of 5-HT receptors is generally organized in seven classes, termed 5-HT₁₋₇. The major classes, 5-HT₁, 5-HT₂, and 5-HT₆, possibly evolved from a primordial, ancestral 5-HT receptor over 750 million years ago and are found in diverse phyla from planarians, nematodes, insects to vertebrates, including man^[61-63]. The 5-HT₅ and 5-HT₇ receptor classes probably deviated from 5-HT₁ 650 to 700 million years ago^[61,62]. Since these events predate the estimated divergence of protostomes and deuterostomes about 600 to 650 million years ago^[66], the invertebrate and vertebrate serotonergic systems are supposed to own roughly the same main receptor classes^[61,67]. However, further gene duplications and structural/functional differentiation led to various subtypes within each main class, and these subtypes are considered to have evolved independently in invertebrates and vertebrates^[61,63].

Insect 5-HT receptors are classified based on sequence similarities with their counterparts in vertebrates. Within the same receptor class, the signaling properties seem very well maintained between insects and vertebrates, but their pharmacological characteristics vary significantly. Compared to their mammalian (and other vertebrate) counterparts, relatively few data about pharmacological properties of insect 5-HT receptors are available.

Table 1: The vertebrate 5-HT receptor types, their G protein coupling characteristics and primary responses. Receptor subfamilies with representatives in insects are indicated (•). Abbreviations: AC, adenylyl cyclase; PLC, phospholipase C

Family	G-protein	Response
5-HT ₁ •	G _{i/o}	AC inhibition
5-HT ₂ •	G _{q/11}	PLC activation
5-HT ₃		Ligand-gated ion channel
5-HT ₄	G _s	AC activation
5-HT ₅	G _{i/o}	AC inhibition
5-HT ₆	G _s	AC activation
5-HT ₇ •	G _s	AC activation

A recent study provides evidence that agonists of 5-HT receptors may constitute potential lead compounds for new insecticide discovery^[68] (see § 6.2).

5-HT receptors in vertebrates

Different classification schemes for 5-HT receptors have been used for more than 50 years. Initially, Gaddum and Picarelli (1957)^[69] described D- and M-type receptors, based on pharmacological studies with dibenzylamine and morphine. In radio-ligand binding studies on brain homogenates of rat, Peroutka and Snyder (1979)^[70] demonstrated the presence of two distinct 5-HT receptor binding sites, named 5-HT₁ and 5-HT₂. These data were amalgamated whereby three main groups of 5-HT receptors were considered, namely 5-HT₁, 5-HT₂ and 5-HT₃ (the latter two corresponding to the D and M receptors, respectively)^[71]. This nomenclature still forms the basis for the current classification of 5-HT receptors.

Presently, vertebrate 5-HT receptors are divided into seven main 5-HT receptor families (termed 5-HT₁₋₇) based on their gene organization, sequence similarities, pharmacological properties and downstream signaling pathways^[14,64,65,72-74]. Six of these belong to the rhodopsin-like family of GPCRs and the lone exception, 5-HT₃, is a cation-selective chloride-channel that belongs to the ligand-gated ion-channel superfamily. This receptor is composed of five pseudo-symmetrical subunits that surround the central ion-channel. Each subunit contains an extracellular part with a ligand-binding domain, a TM with four α -helices and a cytoplasmatic domain. Activation of the receptor, by binding of 5-HT, leads to the opening of the channel and a quick inward ion current that will cause a depolarization of the plasma membrane^[14,65,75]. 5-HT₁ and 5-HT₅ couple to G_{i/o}-proteins, which inhibit adenylyl cyclase activity and thus decrease the intracellular cyclic AMP (cAMP) levels. 5-HT₂ couples to G_{q/11}-proteins that activate phospholipase C (PLC). PLC hydrolyses membrane bound phosphoinositides, like phosphatidyl inositol

4,5-bisphosphate (PIP₂). This results in the formation of diacylglycerol (DAG) and inositol phosphates like inositol 1,4,5-trisphosphate (IP₃). IP₃ causes Ca²⁺-exemption in the cell after binding to ligand-regulated Ca²⁺-channels on the membrane of intracellular Ca²⁺-storage compartments, such as the endoplasmic reticulum. DAG remains associated to the cell membrane where it will activate, together with the higher intracellular Ca²⁺-concentration, Ca²⁺-dependent protein kinase (PKC). PKC phosphorylates serine and threonine residues of protein substrates, which alters the functional properties of these proteins. 5-HT₄, 5-HT₆ and 5-HT₇ couple to G_s-proteins; the G_{as}-subunit stimulates adenylyl cyclase activity and hence increases the intracellular cAMP levels (Table 1)^[14,63-65,73,76]. The main receptor families contain different subtypes and isoforms (designated a, b, c ...). But even this amount of physical diversity underscores the physiological relevance of 5-HT. An even greater degree of operational diversity can be expected, based on the existence of a great number of splice and RNA editing variants for several 5-HT receptors (see § 5.1.1). In addition, they can be modulated by accessory proteins and chaperones, and can form homo- or heteromers both at the GPCR and the ligand-gated channel level (for a review, see: Hannon and Hoyer, 2008^[65]).

5-HT receptor multiplicity in vertebrates

There are five vertebrate 5-HT₁ receptor subtypes, all encoded by intronless genes: 5-HT_{1A,B,D,E,F}^[77-86]. In addition, two polymorphisms have been found to alter the extracellular amino-terminal region of the human 5-HT_{1A} receptor; one is associated with Tourette's syndrome, and one results in loss of response to 5-HT^[87,88]. The current 5-HT receptor nomenclature does not contain a 5-HT_{1C} receptor. This receptor was formerly (incorrectly) classified in the 5-HT₁ receptor family, but was renamed 5-HT_{2C}^[74]. This 5-HT_{2C} receptor was the first cloned receptor of the 5-HT₂ receptor family^[89,90] which now contains three subtypes: 5-HT_{2A-C}^[91-94]. The 5-HT_{2C} receptor forms different isoforms through RNA editing by adenosine deaminases, which affects agonist potency, activation of PLC and selectivity of G protein coupling^[95-97]. In mammals, a third family of 5-HT GPCRs comprises at least nine 5-HT₄ isoforms (5-HT_{4a-g}, 5-HT_{4i} and 5-HT_{4n}) resulting from alternative splicing starting from a conserved leucine residue in the intracellular C-terminus. Another variant, 5-HT_{4hb} contains a 14 residue insertion in the second extracellular loop compared to the 5-HT_{4b} variant^[98-106]. Moreover, different isoforms that are co-expressed in the same tissue were shown to form dimers^[107]. Two subtypes of 5-HT₅ receptors (5-HT_{5A} and 5-HT_{5B}) have been cloned in rodents, sharing 70% overall sequence identity^[108-111]. In humans, the 5-HT_{5A} constitutes the only functional 5-HT₅ receptor. The 5-HT_{5B} receptor gene has been mapped

to the human genome, but does not encode a functional protein^[112,113]. The first functional 5-HT₆ receptor was cloned from rat cDNA starting from the sequence of the rat histamine H₂ receptor^[114]. The rat sequence was later corrected, together with cloning of the human 5-HT₆ receptor sequence^[115]. A degenerate PCR approach on human cDNA was used to identify a nonfunctional 5-HT₆ receptor splice variant^[116]. Members of the 5-HT₇ receptor family were cloned around the same time from rodent and human. Two introns are found in the mammalian receptor encoding gene, from which one is alternatively spliced resulting in differences in the intracellular C-terminus^[117-125]. In rat and mouse tissue, three 5-HT₇ isoforms were found, called 5-HT_{7a}, 5-HT_{7b} and 5-HT_{7c}. Two human splice variants correspond to isoforms a and b, but a third constitutes a distinct isoform, called 5-HT_{7d}^[118,119,126,127].

5-HT receptors in insects

For many years, the work on 5-HT receptors of invertebrate physiologists and pharmacologists has paralleled that of vertebrate researchers. This resulted in a distinct nomenclature for invertebrate 5-HT receptors that became increasingly confusing when compared to the vertebrate classification. Tierney (2001)^[63] proposed a classification system for invertebrate 5-HT receptors based on the vertebrate nomenclature, considering the common ancestry of insect and vertebrate 5-HT receptors. Indeed, as mentioned above, the main receptor classes are believed to have diverged from a primordial 5-HT receptor before the estimated separation of protostomes and deuterostomes^[61,62,66].

Based on the conserved amino acid sequences and activated second-messenger systems, insect 5-HT receptors can be classified as 5-HT₁, 5-HT₂ and 5-HT₇ type GPCRs. *D. melanogaster* was the first insect species in which four different 5-HT receptors were pharmacologically characterized, namely a 5-HT_{1α/A}, 5-HT_{1β/B}, 5-HT_{2α/A} and 5-HT₇ receptor^[128,129]. In addition, the existence of a second 5-HT₂ receptor (5-HT_{2β/B}) was suggested and recently confirmed experimentally^[41,130,131]. In the field cricket, partial sequences of two 5-HT₁, two 5-HT₂ and a 5-HT₇ receptor were identified as well^[132,133]. Two 5-HT₁ receptors were also predicted in the genome of *T. castaneum*, one of these was characterized^[134,135] and two 5-HT₁ splice variants were described in *Papilio xuthus*^[136]. On the contrary, in the locust *L. migratoria* only one 5-HT₁ receptor was partially cloned and also in *A. mellifera* and *P. americana* only one 5-HT₁ receptor was characterized thus far^[137-139]. As in *D. melanogaster* and *G. bimaculatus*, two 5-HT₂ receptors were identified from *A. mellifera*^[41,128,130-133,140], while in other insects, such as *T. castaneum* and the locust *L. migratoria*,

only one 5-HT₂ receptor has been found thus far^[134,137]. Lastly, ever since the first insect 5-HT₇ receptor was identified from a *Drosophila* genomic and a cDNA library^[129,141], (partial) cDNA sequences of 5-HT₇ receptors from other insects were cloned and some of these were functionally characterized^[133,134,137,142,143]. Based on the data from characterized insect 5-HT receptors, most insects seem to possess one 5-HT₇ type serotonin receptor, but may possess two 5-HT₁ and/or two 5-HT₂ receptors.

Similarly to vertebrate receptors, alternative splicing seems a general mechanism for creating extra multiplicity in the insect 5-HT receptor system (for example the 5-HT₁ receptors of *D. melanogaster* and *P. xuthus*)^[129,136].

Insect 5-HT receptors often show high expression in nervous tissue (e.g. brain and ventral nerve cord)^[36,37,133,135,138,142-144]. High 5-HT₁ protein levels in the central nervous system were associated with involvement in visual information processing in honey bee foragers^[138] and possible roles in neuroendocrine secretion processes and motility of the gut in cockroaches^[139]. The expression of 5-HT₇ in different parts of the honey bee brain suggested a role for the receptor in neural pathways for information processing, learning and memory^[142]. Several insect 5-HT₇ receptors were also found to be highly expressed in the salivary gland (FlyAtlas, <http://www.flyatlas.cor/>)^[133,145] and suggested to be involved in 5-HT induced saliva secretion, which was shown to be mediated via an increase in intracellular cAMP in several insect species^[44,45,139,146-149]. Immunohistochemistry with antibodies against the *A. aegypti* 5-HT₇ receptor labeled two axons that run in parallel along the hindgut^[144]. In *L. migratoria*, high serotonergic immunoreactivity was measured in the midgut and application of 5-HT induced relaxation of the midgut circular muscle^[150]. 5-HT₇ receptor-mediated regulation of visceral muscle activity in the gastrointestinal tract may be evolutionary conserved between invertebrates and vertebrates, since also mammalian 5-HT₇ receptors were demonstrated to mediate smooth muscle relaxation of the gastrointestinal tract^[151].

Pharmacology of vertebrate 5-HT receptors

The pharmacology of mammalian 5-HT receptor subtypes has been extensively studied (<http://www.iuphar-db.org>). We will only briefly discuss a selection of agonists and antagonists, focusing on 5-HT₁, 5-HT₂ and 5-HT₇ receptors, since orthologs of these receptor types were found in insects.

5-HT_{1A} receptors are probably the most extensively examined receptor group. One of the first major full and selective 5-HT_{1A} receptor agonists to be discovered was

Table 2. Overview of pharmacological agents tested on insect serotonin receptors

Receptor	Transduction	Agonists	EC ₅₀	Antagonists	Cells	References
<i>Drome</i> 5-HT1A CAA77570	decrease cAMP increase IP	5-HT 8-OH-DPAT		dihydroergocryptine prazosin d-butaclamol methysergide l-butaclamol yohimbine methiothepin	Cos-7	Saudou <i>et al.</i> , 1992 ^[129]
		5-HT	1 μM		HEK293T	Gasque <i>et al.</i> , 2013 ^[41]
<i>Drome</i> 5-HT1B CAA77571	decrease cAMP increase IP	5-HT 8-OH-DPAT		dihydroergocryptine d-butaclamol prazosin methysergide yohimbine l-butaclamol methiothepin	Cos-7	Saudou <i>et al.</i> , 1992 ^[129]
		5-HT	625 nM		HEK293T	Gasque <i>et al.</i> , 2013 ^[41]
<i>Apime</i> 5-HT1A CBI75449	decrease cAMP	5-HT 5-CT 5-MT	16.9 nM 700.5 nM 3.63 μM	methiothepin prazosin (partial) WAY-100635 (partial)	HEK293	Thamm <i>et al.</i> , 2010 ^[138]
<i>Trica</i> 5-HT1 KC196076		5-HT α-Me5-HT 5-CT 5-MT 8-OH-DPAT	95 nM 10.7 μM 24.7 μM 91.8 μM 551.0 μM	prazosin methiothepin methysergide d-butaclamol SB-269970 WAY-100635 <i>ketanserine</i> (NR) <i>mianserine</i> (NR) <i>yohimbine</i> (NR)	CHO-WTA11	Vleugels <i>et al.</i> , 2013 ^[135]
	decrease cAMP	5-HT	82.7 nM		HEK293	Vleugels <i>et al.</i> , 2013 ^[135]
<i>Peram</i> 5-HT1 FN298392	decrease cAMP	5-HT 5-MT	130 nM (partial)	WAY-100635 (inverse) Methiothepin (neutral)	HEK293	Troppmann <i>et al.</i> , 2010 ^[139]
<i>Drome</i> 5-HT2(A) CAA57429	?	5-HT N-acetyl-5-HT α-Me5-HT 2-Me5-HT tryptamine 1-Me5-HT 5-MT quipazine 5-MT 8-OH-DPAT 5-CT		ritanserine ketanserine pizotifen setoperone spiperone cyproheptadine TFMPP mesulergine methysergide methiothepine rauwolscine buspirone yohimbine bufotenine sulpiride mianserine clozapine cis-flupenthixol haloperidol methiothepin	Cos-1	Colas <i>et al.</i> , 1995 ^[128]
		5-HT	99 nM		HEK293T	Gasque <i>et al.</i> , 2013 ^[41]
<i>Drome</i> 5-HT2B CG42796		5-HT	293 nM	methiothepin	HEK293T	Gasque <i>et al.</i> , 2013 ^[41]
<i>Apime</i> 5-HT2A FR727107	increase Ca2+	5-HT 5-MT 8-OH-DPAT	25.7 nM 70 nM 55.9 μM	SB-269970 mianserine cyproheptadine methiothepin clozapine methysergide	HEK293	Thamm <i>et al.</i> , 2013 ^[140]
<i>Apime</i> 5-HT2B FR727108	increase Ca2+	5-HT 5-MT 8-OH-DPAT	32.5 nM 60.4 nM 561.5 nM	cyproheptadine ketanserine mianserine clozapine	HEK293	Thamm <i>et al.</i> , 2013 ^[140]
<i>Drome</i> 5-HT7 M55533	increase cAMP	5-HT 2-Me5HT 5-MT d-LSD	60 nM 0.6 μM 0.8 μM 1.5 μM	dihydroergocryptine d-butaclamol methysergide	NIH 3T3	Witz <i>et al.</i> , 1990 ^[141]

	increase cAMP	8-OH-DPAT 5-HT 8-OH-DPAT	NR	dihydroergocryptine d-butaclamol methysergide prazosin l-butaclamol yohimbine metitepine	Cos-7	Saudou <i>et al.</i> , 1992 ^[129]
		5-HT	12 nM		HEK293T	Gasque <i>et al.</i> , 2013 ^[41]
<i>Apime</i> 5-HT7 CAJ28210	increase cAMP	5-HT 5-CT AS-19 8-OH-DPAT	1.1-1.8 nM 24-40 nM moderate poor	methiothepin (inverse) SB-269970 (poor) <i>clozapine</i> (NR)	HEK293	Schlenstedt <i>et al.</i> , 2006 ^[142]
<i>Trica</i> 5-HT7 XP_966577		5-HT α -Me5-HT 5-CT 5-MT 8-OH-DPAT	27.3 nM 5.2 μ M 22.2 μ M 30.3 μ M 38.8 μ M	ketanserin methysergide methiothepine d-butaclamol prazosin SB-269970 yohimbine WAY-100635 mianserin	CHO-WTA11	Vleugels <i>et al.</i> , 2014 ^[143]
	increase cAMP	5-HT	6.6 nM		HEK293	Vleugels <i>et al.</i> , 2014 ^[143]
<i>Aedae</i> 5-HT7 AAG49292	increase cAMP	5-HT 5-CT 8-OH-DPAT pimozide	39.5 nM moderate moderate poor		CHO-K1	Lee and Pietrantonio, 2003 ^[169]

Pharmacological profiles of heterologously expressed insect 5-HT receptors. Agonists and antagonists are shown according to decreasing potency (per receptor and cell type). Agonists and antagonists that exhibited no response (NR) are shown in grey and italic. Abbreviations used: 5-CT, 5-carboxamidotryptamine; 5-HT, 5-hydroxytryptamine; 5-MT, 5-methoxytryptamine; 8-OH-DPAT, 8-hydroxy-2-(di-n-propylamino)tetralin; *Aedae*, *Aedes aegypti*; *Apime*, *Apis mellifera*; AS-19, (2S)-N,N-dimethyl-5-(1,3,5-trimethylpyrazol-4-yl)-1,2,3,4-tetrahydronaphthalen-2-amine; cAMP, cyclic adenosine monophosphate; CHO, Chinese hamster ovary; Cos, fibroblast-like cell line derived from monkey kidney tissue; *Drome*, *Drosophila melanogaster*; EC₅₀, half maximal effective concentration; HEK, human embryonic kidney; IP, inositol phosphate; Me5-HT, methyl hydroxytryptamine; NIH, fibroblast cell line derived from mouse embryos; *Peram*, *Periplaneta americana*; SB-269970, 2R)-1-[(3-hydroxyphenyl)sulfonyl]-2-(2-(4-methyl-1-piperidiny)ethyl)pyrrolidine; *Trica*, *Tribolium castaneum*; WAY-100635, N-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-N-(2-pyridyl)cyclohexanecarboxamide.

8-OH-DPAT (8-hydroxy-2-(di-n-propylamino)tetralin)^[84,152]. It was later shown to have some minor affinity for 5-HT_{1D} and 5-HT₇ receptors as well (see IUPHAR database and references therein). Another well-known agonist of 5-HT₁ and 5-HT₇ receptors is 5-CT (5-carboxamidotryptamine)^[117,153,154]. Agonists with minor affinity for 5-HT₁ receptors are 5-MeOT (5-methoxytryptamine) which has high agonistic activity on 5-HT_{2A} and 5-HT₇ receptors, and α m-5-HT (α -methylserotonin) which has higher specificity for 5-HT₂ receptors compared to all other 5-HT receptor (sub)types^[117,153,155-157]. Antagonists with very high specificity were found for some mammalian 5-HT receptors. The piperazine drug WAY-100635 is a potent antagonist of 5-HT_{1A} receptors^[158,159], while SB-269970 is a potent antagonist of 5-HT₇ receptors^[160,161]. However, WAY-100635 was also shown to be a D₄ dopamine receptor agonist^[162] and SB-269970 was able to block α 2-adrenoceptors to some extent^[163]. Several potent 5-HT₂ receptor antagonists with only minor binding affinity for other 5-HT receptors are known, including mianserin, ketanserin, and butaclamol (see IUPHAR database and references therein). However, all have affinity for other biogenic amine receptors as well, with a particularly high binding affinity of ketanserin for

α 1-adrenergic receptors^[164] and of butaclamol for dopamine receptors^[165-168]. Several non-selective 5-HT receptor antagonists were characterized, such as methiothepin and methysergide.

Pharmacology of insect 5-HT receptors

Despite the sequence and signaling similarities, the pharmacology of insect 5-HT receptors is curiously different from that of their vertebrate counterparts. The long evolutionary history of the 5-HT signaling systems after the divergence of protostomes and deuterostomes, allowed for further differentiation into subtypes independently in insects and vertebrates^[63,64]. During this evolution, selection was likely based on functionally important receptor characteristics, such as ligand binding and G protein coupling, and not the conservation of recognition sites for man-made, synthetic ligands^[143].

Although the pharmacology of the mammalian receptors is thoroughly studied (and drugs affecting biogenic amine receptor binding are among the most important in the armory of modern pharmacology), the specificity of these drugs towards insect receptors has never been comprehensively tested and profound characterization studies of insect 5-HT

receptors are scarce. Consequently much less is known about the pharmacology of these receptors. However, it is a recurring fact that the synthetic agonists, designed for vertebrate receptors, are much less potent and often also less efficient than the natural ligand 5-HT on insect 5-HT receptors. For example, the mammalian 5-HT_{1/7} receptor agonists, 8-OH-DPAT and 5-CT, act only as poor agonists on insect 5-HT_{1/7} receptors^[135,138,139,142,143,169]. Little information is also available about selective or potent antagonists of insect 5-HT receptors and only limited similarity is observed with the mammalian receptor pharmacology (Table 2). The potent and selective mammalian 5-HT_{1A} antagonist WAY-100635 showed only minor antagonistic activity on insect 5-HT₁ receptors, i.e. from *A. mellifera*, *P. americana* and *T. castaneum*^[135,138,139]. The mammalian 5-HT₇ receptor antagonist SB-269970 has been shown to be a potent antagonist of the 5-HT₇ receptor from *C. vicina*, but had only poor influence on activity of the 5-HT₇ receptors from *A. mellifera* and *T. castaneum*^[142,143,145]. The non-selective 5-HT antagonists of vertebrate 5-HT receptors, methysergide and methiothepin, were also able to decrease insect 5-HT receptor activity^[135,143].

5-HT receptors as pharmacological targets

5-HT receptors as drug targets

Depression and anxiety disorders are two well-known health problems for which compounds that influence 5-HT receptor activity are prescribed. About 80% of all antidepressants on the market are selective serotonin re-uptake inhibitors (SSRIs), preventing 5-HT uptake from the synapse by SERTs. A main disadvantage of SSRIs is that the onset of their therapeutic action requires weeks of treatment. The use of partial agonists of 5-HT_{1A} receptors and inhibitors of 5-HT_{2A} receptors was shown to accelerate clinical antidepressant effects^[23,170-173]. Recently, two new antidepressant drugs (vilazodone and vortioxetine) which combine partial 5-HT_{1A} agonist properties with SERT blockade are approved by the FDA (U.S. Food and Drug Administration) to treat depression (Reuters Press Release, 2011; U.S. Food and Drug Administration Press Release, 2013). Partial 5-HT₁ agonists, including certain azapirones, are also used as anxiolytic drugs^[172,174]. It is very likely that 5-HT₄ receptor partial agonists could behave as rapid and effective antidepressants as well^[175].

The most widely used anti-migraine drugs are triptans which bind with high affinity to 5-HT_{1B/D/F} receptors. However, 5-HT_{1F} is the most attractive target for new anti-migraine drugs since activation of 5-HT_{1B} induces cardiac adverse effects and 5-HT_{1D} agonists – although effective in animal models of migraine – failed to properly

attenuate migraine attacks in clinical trials. One selective 5-HT_{1F} agonist (lasmiditan) has successfully completed phase 2 clinical trials (in 2007 for an intravenous and in 2010 for an oral application)^[176]. Also agonists of other 5-HT receptors are used in treatment of various health problems (for example^[177,178]).

5-HT receptors as potential insecticide targets

The poor affinity and selectivity on insect receptors of the synthetic compounds that were originally designed for mammalian receptor subtypes indicate the importance to look for pharmacological agents that target the insect receptor types more specifically, both for research and insect control. Indeed, although 5-HT receptors are involved in many key processes of insect life, no insecticides that target this receptor group are available at this moment. However, one study provided evidence that 5-HT receptor agonists may form potential lead compounds for new insecticide discovery^[68]. They used the 5-HT_{1A} agonist PAPP (1-[(4-aminophenyl)ethyl]-4-[3-(trifluoromethyl)phenyl]piperazine) as a lead compound for new insecticides with novel mode of action. The PAPP scaffold was used to design and synthesize a series of compounds that were evaluated for biological activity against the lepidopteran armyworm *Pseudaletia separata*. Most of the compounds indeed displayed growth-inhibiting potential or induced larval death. Another study in *Drosophila* shows that the general 5-HT receptor antagonist methiothepin inhibits feeding through 5-HT_{2A}^[41]. Given the ubiquity of 5-HT signaling in the animal kingdom, any insect control agent targeting 5-HT receptors would have to be thoroughly tested for specificity to avoid toxicity on a wide range of species^[55,179].

Concluding remarks

Several interesting studies have highlighted the omnipresent and overwhelming functions of serotonin both in vertebrates as in invertebrates. The number of studies on the role of serotonin in insects is still relatively limited, but these studies already underline the immense importance of serotonin signaling in nearly all major physiological processes in insects.

Identification and classification of insect 5-HT receptors is merely based on sequence similarity with mammalian receptors. Although this information can provide indications for receptor classification and comparison, it does not allow for accurate predictions of the pharmacological characteristics of these receptors. The current pharmacological data available on insect 5-HT receptors suggest that it should be possible to find selective synthetic ligands for specific insect 5-HT receptors. However, a lot of

additional research, contributing both to proper insect 5-HT receptor classification and to understanding their *in vivo* functionality, will be necessary before a selective insecticide can be developed.

Acknowledgements

RV was supported by a PhD fellowship from IWT (Agency for Innovation by Science and Technology in Flanders). HV was supported by the Research Foundation of Flanders (FWO Flanders). Our research is also sponsored by the Interuniversity Attraction Poles program (Belgian Science Policy Grant P7/40) and the KU Leuven Research Foundation (GOA/11/02).

Authors' contributions: RV and HV contributed equally to the manuscript; JV is the senior academic author and revised the manuscript critically.

List of abbreviations

5-CT: 5-carboxamidotryptamine; 5-HT: 5-hydroxytryptamine or serotonin; 5-HTTP: 5-hydroxytryptophan; 5-MeOT: 5-methoxytryptamine; 8-OH-DPAT: 8-hydroxy-2-(di-n-propylamino) tetralin; α M-5-HT: α -methylserotonin; GPCR: G protein-coupled receptor; IP₃: inositol 1,4,5-trisphosphate; DAG: diacylglycerol; DOPA: 3,4-dihydroxyphenylalanine; LSD: lysergide acid diethylamide; PAPP: 1-[(4-aminophenyl)ethyl]-4-[3-(trifluoromethyl)phenyl]piperazine; PKC: Ca²⁺-dependent protein kinase; PLC: phospholipase C; PIP₂: phosphatidyl inositol 4,5-bisphosphate; SERT: SERotonin Transporter; SSRI: selective serotonin re-uptake inhibitor; TM: transmembrane segments; TPH: tryptophan hydroxylase.

References

- Blenau W, Baumann A. Molecular and pharmacological properties of insect biogenic amine receptors: lessons from *Drosophila melanogaster* and *Apis mellifera*. Arch Insect Biochem Physiol 2001; 48:13-38.
- Pflüger HJ, Stevenson PA. Evolutionary aspects of octopaminergic systems with emphasis on arthropods. Arthropod Struct Dev 2005; 34:379-396.
- Robertson HA, Juorio AV. Octopamine and some related noncatecholic amines in invertebrate nervous systems. Int Rev Neurobiol 1976; 19:173-224.
- Roeder T, Degen J, Dyczkowski C, Gewecke M. Pharmacology and molecular biology of octopamine receptors from different insect species. Prog Brain Res 1995; 106:249-258.
- Juorio AV, Robertson HA. Identification and distribution of some monoamines in tissues of the sunflower star, *Pycnopodia helianthoides* (Echinodermata). J Neurochem 1977; 28:573-579.
- Moret F, Guiland JC, Coudouel S, Rochette L, Vernier P. Distribution of tyrosine hydroxylase, dopamine, and serotonin in the central nervous system of amphioxus (*Branchiostoma lanceolatum*): implications for the evolution of catecholamine systems in vertebrates. J Comp Neurol 2004; 468:135-150.
- Premont RT, Gainetdinov RR, Caron MG. Following the trace of elusive amines. Proc Natl Acad Sci U S A 2001; 98:9474-9475.
- Roberts KM, Fitzpatrick PF. Mechanisms of tryptophan and tyrosine hydroxylase. IUBMB Life 2013; 65:350-357.
- Fitzpatrick PF. Tetrahydropterin-dependent amino acid hydroxylases. Annu Rev Biochem 1999; 68:355-381.
- Hufton SE, Jennings IG, Cotton RG. Structure and function of the aromatic amino acid hydroxylases. Biochem J 1995; 311:353-366.
- Kim-Ha J, Smith JL, Macdonald PM. oskar mRNA is localized to the posterior pole of the *Drosophila* oocyte. Cell 1991; 66:23-35.
- Lovenberg W, Weissbach H, Udenfriend S. Aromatic L-amino acid decarboxylase. J Biol Chem 1962; 237:89-93.
- Donly BC, Caveney S. A transporter for phenolamine uptake in the arthropod CNS. Arch Insect Biochem Physiol 2005; 59:172-183.
- Nichols DE, Nichols CD. Serotonin receptors. Chem Rev 2008; 108:1614-1641.
- Scavone C, McKee M, Nathanson JA. Monoamine uptake in insect synaptosomal preparations. Insect Biochem Mol Biol 1994; 24:589-597.
- Torres GE, Carneiro A, Seamans K, Fiorentini C, Sweeney A, Yao WD, et al. Oligomerization and trafficking of the human dopamine transporter. Mutational analysis identifies critical domains important for the functional expression of the transporter. J Biol Chem 2003; 278:2731-2739.
- Corey JL, Quick MW, Davidson N, Lester HA, Guastella J. A cocaine-sensitive *Drosophila* serotonin transporter: cloning, expression, and electrophysiological characterization. Proc Natl Acad Sci U S A 1994; 91:1188-1192.
- Demchyshyn LL, Pristupa ZB, Sugamori KS, Barker EL, Blakely RD, Wolfgang WJ, et al. Cloning, expression, and localization of a chloride-facilitated, cocaine-sensitive serotonin transporter from *Drosophila melanogaster*. Proc Natl Acad Sci U S A 1994; 91:5158-5162.
- Porzgen P, Park SK, Hirsh J, Sonders MS, Amara SG. The antidepressant-sensitive dopamine transporter in *Drosophila melanogaster*: a primordial carrier for catecholamines. Mol Pharmacol 2001; 59:83-95.
- Malutan T, McLean H, Caveney S, Donly C. A high-affinity octopamine transporter cloned from the central nervous system of cabbage looper *Trichoplusia ni*. Insect Biochem Mol Biol 2002; 32:343-357.
- Nathanson JA, Hunnicutt EJ, Kantham L, Scavone C. Cocaine as a naturally occurring insecticide. Proc Natl Acad Sci U S A 1993; 90:9645-9648.
- Roeder T, Gewecke M. Octopamine uptake systems in thoracic ganglia and leg muscles of *Locusta migratoria*. Comp Biochem Physiol 1989; 94:143-147.
- Jones BJ, Blackburn TP. The medical benefit of 5-HT research. Pharmacol Biochem Behav 2002; 71:555-568.
- Dewhurst SA, Croker SG, Ikeda K, McCaman RE. Metabolism of

- biogenic amines in *Drosophila* nervous tissue. Comp Biochem Physiol B 1972; 43:975-981.
25. Fickbohm DJ, Spitzer N, Katz PS. Pharmacological manipulation of serotonin levels in the nervous system of the opisthobranch mollusc *Tritonia diomedea*. Biol Bull 2005; 209:67-74.
26. Sloley BD. Metabolism of monoamines in invertebrates: the relative importance of monoamine oxidase in different phyla. Neurotoxicology 2004; 25:175-183.
27. Rapport MM, Green AA, Page IH. Serum vasoconstrictor, serotonin; isolation and characterization. J Biol Chem 1948; 176:1243-1251.
28. Rapport MM, Green AA, Page IH. Crystalline serotonin. Science 1948; 108:329-330.
29. Rapport MM. Serum vasoconstrictor (serotonin) the presence of creatinine in the complex; a proposed structure of the vasoconstrictor principle. J Biol Chem 1949; 180:961-969.
30. Erspamer V, Asero B. Identification of enteramine, the specific hormone of the enterochromaffin cell system, as 5-hydroxytryptamine. Nature 1952; 169:800-801.
31. Vialli M, Erspamer V. Ricerche sul secreto delle cellule enterocromaffini. Zellforschung mikr Anatomie 1937; 27:81-99.
32. Berger M, Gray JA, Roth BL. The expanded biology of serotonin. Annu Rev Med 2009; 60:355-366.
33. Gershon MD, Tack J. The serotonin signaling system: From basic understanding to drug development-for functional GI disorders. Gastroenterology 2007; 132:397-414.
34. Twarog BM, Page IH. Serotonin content of some mammalian tissues and urine and a method for its determination. Am J Physiol 1953; 175:157-161.
35. Woolley DW, Shaw E. Some neurophysiological aspects of serotonin. Br Med J 1954; 2:122-126.
36. Yuan Q, Joiner WJ, Sehgal A. A sleep-promoting role for the *Drosophila* serotonin receptor 1A. Curr Biol 2006; 16:1051-1062.
37. Yuan Q, Lin F, Zheng X, Sehgal A. Serotonin modulates circadian entrainment in *Drosophila*. Neuron 2005; 47:115-127.
38. Neckameyer WS. A trophic role for serotonin in the development of a simple feeding circuit. Dev Neurosci 2010; 32:217-237.
39. Dacks AM, Nickel T, Mitchell BK. An examination of serotonin and feeding in the flesh fly *Neobellieria bullata* (Sarcophagidae: Diptera). J Insect Behav 2003; 16:1-21.
40. French AS, Simcock KL, Rolke D, Gartside SE, Blenau W, Wright GA. The role of serotonin in feeding and gut contractions in the honeybee. J Insect Physiol 2014; 61:8-15.
41. Gasque G, Conway S, Huang J, Rao Y, Vosshall LB. Small molecule drug screening in *Drosophila* identifies the 5HT2A receptor as a feeding modulation target. Sci Rep 2013; 3:srep02120.
42. Novak MG, Ribeiro JMC, Hildebrand JG. 5-Hydroxytryptamine in the salivary-glands of adult female *Aedes aegypti* and its role in regulation of salivation. J Exp Biol 1995; 198:167-174.
43. Novak MG, Rowley WA. Serotonin depletion affects blood-feeding but not host-seeking ability in *Aedes triseriatus* (Diptera: Culicidae). J Med Entomol 1994; 31:600-606.
44. Berridge MJ, Patel NG. Insect salivary glands - Stimulation of fluid secretion by 5-hydroxytryptamine and adenosine-3',5'-monophosphate. Science 1968; 162:462-463.
45. Walz B, Baumann O, Krach C, Baumann A, Blenau W. The aminergic control of cockroach salivary glands. Arch Insect Biochem Physiol 2006; 62:141-152.
46. Donini A, O'Donnell MJ, Orchard I. Differential actions of diuretic factors on the Malpighian tubules of *Rhodnius prolixus*. J Exp Biol 2008; 211:42-48.
47. Maddrell SH, Herman WS, Mooney RL, Overton JA. 5-Hydroxytryptamine: a second diuretic hormone in *Rhodnius prolixus*. J Exp Biol 1991; 156:557-566.
48. Kaplan DD, Zimmermann G, Suyama K, Meyer T, Scott MP. A nucleostemin family GTPase, NS3, acts in serotonergic neurons to regulate insulin signaling and control body size. Genes Dev 2008; 22:1877-1893.
49. Ruaud AF, Thummel CS. Serotonin and insulin signaling team up to control growth in *Drosophila*. Genes Dev 2008; 22:1851-1855.
50. Sitaraman D, Laferriere H, Birman S, Zars T. Serotonin is critical for rewarded olfactory short-term memory in *Drosophila*. J Neurogenet 2012; 26:238-244.
51. Sitaraman D, Zars M, Laferriere H, Chen YC, Sable-Smith A, Kitamoto T, et al. Serotonin is necessary for place memory in *Drosophila*. Proc Natl Acad Sci U S A 2008; 105:5579-5584.
52. Bicker G, Menzel R. Chemical codes for the control of behaviour in arthropods. Nature 1989; 337:33-39.
53. Menzel R, Heyne A, Kinzel C, Gerber B, Fiala A. Pharmacological dissociation between the reinforcing, sensitizing, and response-releasing functions of reward in honeybee classical conditioning. Behav Neurosci 1999; 113:744-754.
54. Mercer AR, Menzel R. The effects of biogenic amines on conditioned and unconditioned responses to olfactory stimuli in the honeybee *Apis mellifera*. J Comp Physiol A 1982; 145:363-368.
55. Anstey ML, Rogers SM, Ott SR, Burrows M, Simpson SJ. Serotonin mediates behavioral gregarization underlying swarm formation in desert locusts. Science 2009; 323:627-630.
56. Ott SR, Verlinden H, Rogers SM, Brighton CH, Quah PS, Vleugels RK, et al. Critical role for protein kinase A in the acquisition of gregarious behavior in the desert locust. Proc Natl Acad Sci U S A 2012; 109:E381-E387.
57. Dierick HA, Greenspan RJ. Serotonin and neuropeptide F have opposite modulatory effects on fly aggression. Nat Genet 2007; 39:678-682.
58. Wood SJ, Osborne RH, Cattell KJ, Banner SE. FMRFamide and related peptides modulate the actions of 5-hydroxytryptamine and proctolin on the foregut of the locust, *Schistocerca gregaria*. Biochem Soc Trans 1990; 18:384-385.
59. Evans PD, Myers CM. Peptidergic and aminergic modulation of insect skeletal muscle. J Exp Biol 1986; 124:143-176.
60. Fournier B, Guerineau N, Mollard P, Girardie J. Effects of two neuronal antidiuretic molecules, neuroparsin and 5-hydroxytryptamine, on cytosolic free calcium monitored with indo-1 in epithelial and muscular cells of the African locust rectum. Biochim Biophys Acta 1994; 1220:181-187.

61. Peroutka SJ, Howell TA. The molecular evolution of G protein-coupled receptors: focus on 5-hydroxytryptamine receptors. *Neuropharmacology* 1994; 33:319-324.
62. Walker RJ, Brooks HL, Holden-Dye L. Evolution and overview of classical transmitter molecules and their receptors. *Parasitology* 1996; 113 Suppl:S3-33.
63. Tierney AJ. Structure and function of invertebrate 5-HT receptors: a review. *Comp Biochem Physiol A* 2001; 128:791-804.
64. Hoyer D, Hannon JP, Martin GR. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav* 2002; 71:533-554.
65. Hannon J, Hoyer D. Molecular biology of 5-HT receptors. *Behav Brain Res* 2008; 195:198-213.
66. Ayala FJ, Rzhetsky A, Ayala FJ. Origin of the metazoan phyla: Molecular clocks confirm paleontological estimates. *Proc Natl Acad Sci U S A* 1998; 95:606-611.
67. Hauser F, Cazzamali G, Williamson M, Blenau W, Grimmelhuijzen CJP. A review of neurohormone GPCRs present in the fruitfly *Drosophila melanogaster* and the honey bee *Apis mellifera*. *Progr Neurobiol* 2006; 80:1-19.
68. Cai M, Li Z, Fan F, Huang Q, Shao X, Song G. Design and synthesis of novel insecticides based on the serotonergic ligand 1-[(4-aminophenyl)ethyl]-4-[3-(trifluoromethyl)phenyl]piperazine (PAPP). *J Agric Food Chem* 2010; 58:2624-2629.
69. Gaddum JH, Picarelli ZP. Two kinds of tryptamine receptor. *Br J Pharmacol Chemother* 1957; 12:323-328.
70. Peroutka SJ, Snyder SH. Multiple serotonin receptors: differential binding of [3H]5-hydroxytryptamine, [3H]lysergic acid diethylamide and [3H]spiperidol. *Mol Pharmacol* 1979; 16:687-699.
71. Bradley PB, Engel G, Feniuk W, Fozard JR, Humphrey PP, Middlemiss DN, et al. Proposals for the classification and nomenclature of functional receptors for 5-hydroxytryptamine. *Neuropharmacology* 1986; 25:563-576.
72. Gerhardt CC, Van Heerikhuizen H. Functional characteristics of heterologously expressed 5-HT receptors. *Eur J Pharmacol* 1997; 334:1-23.
73. Hoyer D, Clarke DE, Fozard JR, Hartig PR, Martin GR, Mylecharane EJ, et al. International union of pharmacology classification of receptors for 5-hydroxytryptamine (serotonin). *Pharmacol Rev* 1994; 46:157-203.
74. Humphrey PP, Hartig P, Hoyer D. A proposed new nomenclature for 5-HT receptors. *Trends Pharmacol Sci* 1993; 14:233-236.
75. Hoyer D. Serotonin 5-HT3, 5-HT4, and 5-HT-M receptors. *Neuropsychopharmacology* 1990; 3:371-383.
76. Scheiner R, Baumann A, Blenau W. Aminergic control and modulation of honeybee behaviour. *Curr Neuropharmacol* 2006; 4:259-276.
77. Adham N, Ellerbrock B, Hartig P, Weinshank RL, Branchek T. Receptor reserve masks partial agonist activity of drugs in a cloned rat 5-hydroxytryptamine1B receptor expression system. *Mol Pharmacol* 1993; 43:427-433.
78. Adham N, Kao HT, Schecter LE, Bard J, Olsen M, Urquhart D, et al. Cloning of another human serotonin receptor (5-HT1F): a fifth 5-HT1 receptor subtype coupled to the inhibition of adenylate cyclase. *Proc Natl Acad Sci U S A* 1993; 90:408-412.
79. Amlaiky N, Ramboz S, Boschert U, Plassat JL, Hen R. Isolation of a mouse "5HT1E-like" serotonin receptor expressed predominantly in hippocampus. *J Biol Chem* 1992; 267:19761-19764.
80. Heuring RE, Peroutka SJ. Characterization of a novel 3H-5-hydroxytryptamine binding site subtype in bovine brain membranes. *J Neurosci* 1987; 7:894-903.
81. Hoyer D, Engel G, Kalkman HO. Characterization of the 5-HT1B recognition site in rat brain: binding studies with (-)[125I]iodocyanopindolol. *Eur J Pharmacol* 1985; 118:1-12.
82. Hoyer D, Engel G, Kalkman HO. Molecular pharmacology of 5-HT1 and 5-HT2 recognition sites in rat and pig brain membranes: radioligand binding studies with [3H]5-HT, [3H]8-OH-DPAT, (-)[125I]iodocyanopindolol, [3H]mesulergine and [3H]ketanserin. *Eur J Pharmacol* 1985; 118:13-23.
83. Leonhardt S, Herrick-Davis K, Titeler M. Detection of a novel serotonin receptor subtype (5-HT1E) in human brain: interaction with a GTP-binding protein. *J Neurochem* 1989; 53:465-471.
84. Middlemiss DN, Fozard JR. 8-Hydroxy-2-(di-n-propylamino)-tetralin discriminates between subtypes of the 5-HT1 recognition site. *Eur J Pharmacol* 1983; 90:151-153.
85. Pazos A, Hoyer D, Palacios JM. The binding of serotonergic ligands to the porcine choroid plexus: characterization of a new type of serotonin recognition site. *Eur J Pharmacol* 1984; 106:539-546.
86. Pedigo NW, Yamamura HI, Nelson DL. Discrimination of multiple [3H]5-hydroxytryptamine binding sites by the neuroleptic spiperone in rat brain. *J Neurochem* 1981; 36:220-226.
87. Lam S, Shen Y, Nguyen T, Messier TL, Brann M, Comings D, et al. A serotonin receptor gene (5HT1A) variant found in a Tourette's syndrome patient. *Biochem Biophys Res Commun* 1996; 219:853-858.
88. Nakhai B, Nielsen DA, Linnoila M, Goldman D. Two naturally occurring amino acid substitutions in the human 5-HT1A receptor: glycine 22 to serine 22 and isoleucine 28 to valine 28. *Biochem Biophys Res Commun* 1995; 210:530-536.
89. Hoyer D. Functional correlates of serotonin 5-HT1 recognition sites. *J Recept Res* 1988; 8:59-81.
90. Xie E, Zhu L, Zhao L, Chang LS. The human serotonin 5-HT2C receptor: complete cDNA, genomic structure, and alternatively spliced variant. *Genomics* 1996; 35:551-561.
91. Foguet M, Nguyen H, Le H, Lubbert H. Structure of the mouse 5-HT1C, 5-HT2 and stomach fundus serotonin receptor genes. *Neuroreport* 1992; 3:345-348.
92. Julius D, Huang KN, Livelli TJ, Axel R, Jessell TM. The 5HT2 receptor defines a family of structurally distinct but functionally conserved serotonin receptors. *Proc Natl Acad Sci U S A* 1990; 87:928-932.
93. Kursar JD, Nelson DL, Wainscott DB, Cohen ML, Baez M. Molecular cloning, functional expression, and pharmacological characterization of a novel serotonin receptor (5-hydroxytryptamine2F) from rat stomach fundus. *Mol Pharmacol* 1992; 42:549-557.
94. Pritchett DB, Bach AW, Wozny M, Taleb O, Dal TR, Shih JC, et

- al. Structure and functional expression of cloned rat serotonin 5HT-2 receptor. *EMBO J* 1988; 7:4135-4140.
95. Burns CM, Chu H, Rueter SM, Hutchinson LK, Canton H, Sanders-Bush E, et al. Regulation of serotonin-2C receptor G-protein coupling by RNA editing. *Nature* 1997; 387:303-308.
96. Coupar IM, Desmond PV, Irving HR. Human 5-HT(4) and 5-HT(7) receptor splice variants: are they important? *Curr Neuropharmacol* 2007; 5:224-231.
97. Sanders-Bush E, Fentress H, Hazelwood L. Serotonin 5-HT2 receptors: molecular and genomic diversity. *Mol Interv* 2003; 3:319-330.
98. Bender E, Pindon A, van Oers I, Zhang YB, Gommeren W, Verhasselt P, et al. Structure of the human serotonin 5-HT4 receptor gene and cloning of a novel 5-HT4 splice variant. *J Neurochem* 2000; 74:478-489.
99. Blondel O, Vandecasteele G, Gastineau M, Leclerc S, Dahmoune Y, Langlois M, et al. Molecular and functional characterization of a 5-HT4 receptor cloned from human atrium. *FEBS Lett* 1997; 412:465-474.
100. Bockaert J, Claeysen S, Sebben M, Dumuis A. 5-HT4 receptors: gene, transduction and effects on olfactory memory. *Ann N Y Acad Sci* 1998; 861:1-15.
101. Claeysen S, Faye P, Sebben M, Taviaux S, Bockaert J, Dumuis A. 5-HT4 receptors: cloning and expression of new splice variants. *Ann N Y Acad Sci* 1998; 861:49-56.
102. Claeysen S, Sebben M, Becamel C, Bockaert J, Dumuis A. Novel brain-specific 5-HT4 receptor splice variants show marked constitutive activity: role of the C-terminal intracellular domain. *Mol Pharmacol* 1999; 55:910-920.
103. Gerald C, Adham N, Kao HT, Olsen MA, Laz TM, Schechter LE, et al. The 5-HT4 receptor: molecular cloning and pharmacological characterization of two splice variants. *EMBO J* 1995; 14:2806-2815.
104. Miale J, Berque-Bestel I, Sicsic S, Langlois M, Fischmeister R, Lezoualc'h F. Pharmacological characterization of the human 5-HT(4(d)) receptor splice variant stably expressed in Chinese hamster ovary cells. *Br J Pharmacol* 2000; 131:827-835.
105. Van den Wyngaert I, Gommeren W, Verhasselt P, Jurzak M, Leysen J, Luyten W, et al. Cloning and expression of a human serotonin 5-HT4 receptor cDNA. *J Neurochem* 1997; 69:1810-1819.
106. Vilario MT, Domenech T, Palacios JM, Mengod G. Cloning and characterization of a novel human 5-HT4 receptor variant that lacks the alternatively spliced carboxy terminal exon. RT-PCR distribution in human brain and periphery of multiple 5-HT4 receptor variants. *Neuropharmacology* 2002; 42:60-73.
107. Berthouze M, Ayoub M, Russo O, Rivail L, Sicsic S, Fischmeister R, et al. Constitutive dimerization of human serotonin 5-HT4 receptors in living cells. *FEBS Lett* 2005; 579:2973-2980.
108. Erlander MG, Lovenberg TW, Baron BM, de Lecea L, Danielson PE, Racke M, et al. Two members of a distinct subfamily of 5-hydroxytryptamine receptors differentially expressed in rat brain. *Proc Natl Acad Sci U S A* 1993; 90:3452-3456.
109. Matthes H, Boschert U, Amlaiky N, Grailhe R, Plassat JL, Muscatelli F, et al. Mouse 5-hydroxytryptamine5A and 5-hydroxytryptamine5B receptors define a new family of serotonin receptors: cloning, functional expression, and chromosomal localization. *Mol Pharmacol* 1993; 43:313-319.
110. Plassat JL, Boschert U, Amlaiky N, Hen R. The mouse 5HT5 receptor reveals a remarkable heterogeneity within the 5HT1D receptor family. *EMBO J* 1992; 11:4779-4786.
111. Wisden W, Parker EM, Mahle CD, Grisel DA, Nowak HP, Yocca FD, et al. Cloning and characterization of the rat 5-HT5B receptor. Evidence that the 5-HT5B receptor couples to a G protein in mammalian cell membranes. *FEBS Lett* 1993; 333:25-31.
112. Grailhe R, Grabtree GW, Hen R. Human 5-HT(5) receptors: the 5-HT(5A) receptor is functional but the 5-HT(5B) receptor was lost during mammalian evolution. *Eur J Pharmacol* 2001; 418:157-167.
113. Nelson DL. 5-HT5 receptors. *Curr Drug Targets CNS Neurol Disord* 2004; 3:53-58.
114. Ruat M, Traiffort E, Arrang JM, Tardivel-Lacombe J, Diaz J, Leurs R, et al. A novel rat serotonin (5-HT6) receptor: molecular cloning, localization and stimulation of cAMP accumulation. *Biochem Biophys Res Commun* 1993; 193:268-276.
115. Kohen R, Metcalf MA, Khan N, Druck T, Huebner K, Lachowicz JE, et al. Cloning, characterization, and chromosomal localization of a human 5-HT6 serotonin receptor. *J Neurochem* 1996; 66:47-56.
116. Olsen MA, Nawoschik SP, Schurman BR, Schmitt HL, Burno M, Smith DL, et al. Identification of a human 5-HT6 receptor variant produced by alternative splicing. *Brain Res Mol Brain Res* 1999; 64:255-263.
117. Bard JA, Zgombick J, Adham N, Vaysse P, Branchek TA, Weinshank RL. Cloning of a novel human serotonin receptor (5-HT7) positively linked to adenylate cyclase. *J Biol Chem* 1993; 268:23422-23426.
118. Erdmann J, Nothen MM, Shimron-Abarbanell D, Rietschel M, Albus M, Borrmann M, et al. The human serotonin 7 (5-HT7) receptor gene: genomic organization and systematic mutation screening in schizophrenia and bipolar affective disorder. *Mol Psychiatry* 1996; 1:392-397.
119. Heidmann DE, Metcalf MA, Kohen R, Hamblin MW. Four 5-hydroxytryptamine7 (5-HT7) receptor isoforms in human and rat produced by alternative splicing: species differences due to altered intron-exon organization. *J Neurochem* 1997; 68:1372-1381.
120. Lovenberg TW, Baron BM, Delecea L, Miller JD, Prosser RA, Rea MA, et al. A novel adenylyl cyclase-activating serotonin receptor (5-HT7) implicated in the regulation of mammalian circadian rhythms. *Neuron* 1993; 11:449-458.
121. Meyerhof W, Obermuller F, Fehr S, Richter D. A novel rat serotonin receptor: primary structure, pharmacology, and expression pattern in distinct brain regions. *DNA Cell Biol* 1993; 12:401-409.
122. Plassat JL, Amlaiky N, Hen R. Molecular cloning of a mammalian serotonin receptor that activates adenylate cyclase. *Mol Pharmacol* 1993; 44:229-236.
123. Ruat M, Traiffort E, Leurs R, Tardivellacombe J, Diaz J, Arrang JM, et al. Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT(7)) activating cAMP formation. *Proc Natl Acad Sci U S A* 1993; 90:8547-8551.

124. Shen Y, Monsma FJ, Metcalf MA, Jose PA, Hamblin MW, Sibley DR. Molecular cloning and expression of a 5-hydroxytryptamine 7 serotonin receptor subtype. *J Biol Chem* 1993; 268:18200-18204.
125. Tsou AP, Kosaka A, Bach C, Zuppan P, Yee C, Tom L, et al. Cloning and expression of a 5-hydroxytryptamine7 receptor positively coupled to adenylyl cyclase. *J Neurochem* 1994; 63:456-464.
126. Gellynck E, Laenen K, Andressen KW, Lintermans B, De Martelaere K, Matthys A, et al. Cloning, genomic organization and functionality of 5-HT(7) receptor splice variants from mouse brain. *Gene* 2008; 426:23-31.
127. Heidmann DE, Szot P, Kohen R, Hamblin MW. Function and distribution of three rat 5-hydroxytryptamine7 (5-HT7) receptor isoforms produced by alternative splicing. *Neuropharmacology* 1998; 37:1621-1632.
128. Colas JF, Launay JM, Kellermann O, Rosay P, Maroteaux L. *Drosophila* 5-HT2 serotonin receptor: coexpression with fushi-tarazu during segmentation. *Proc Natl Acad Sci U S A* 1995; 92:5441-5445.
129. Saudou F, Boschert U, Amlaiky N, Plassat JL, Hen R. A family of *Drosophila* serotonin receptors with distinct intracellular signalling properties and expression patterns. *EMBO J* 1992; 11:7-17.
130. Blenau W, Thamm M. Distribution of serotonin (5-HT) and its receptors in the insect brain with focus on the mushroom bodies: lessons from *Drosophila melanogaster* and *Apis mellifera*. *Arthropod Struct Dev* 2011; 40:381-394.
131. Clark AC, del Campo ML, Ewer J. Neuroendocrine control of larval ecdysis behavior in *Drosophila*: complex regulation by partially redundant neuropeptides. *J Neurosci* 2004; 24:4283-4292.
132. Watanabe T, Aonuma H. Identification and expression analyses of a novel serotonin receptor gene, 5-HT2beta, in the field cricket, *Gryllus bimaculatus*. *Acta Biol Hung* 2012; 63 Suppl 2:58-62.
133. Watanabe T, Sadamoto H, Aonuma H. Identification and expression analysis of the genes involved in serotonin biosynthesis and transduction in the field cricket *Gryllus bimaculatus*. *Insect Mol Biol* 2011; 20:619-635.
134. Hauser F, Cazzamali G, Williamson M, Park Y, Li B, Tanaka Y, Predel R, et al. A genome-wide inventory of neurohormone GPCRs in the red flour beetle *Tribolium castaneum*. *Front Neuroendocrinol* 2008; 29:142-165.
135. Vleugels R, Lenaerts C, Baumann A, Vanden Broeck J, Verlinden H. Pharmacological characterization of a 5-HT1-type serotonin receptor in the red flour beetle, *Tribolium castaneum*. *PLoS ONE* 2013; 8:e65052.
136. Ono H, Yoshikawa H. Identification of amine receptors from a swallowtail butterfly, *Papilio xuthus* L.: cloning and mRNA localization in foreleg chemosensory organ for recognition of host plants. *Insect Biochem Mol Biol* 2004; 34:1247-1256.
137. Guo X, Ma Z, Kang L. Serotonin enhances solitariness in phase transition of the migratory locust. *Front Behav Neurosci* 2013; 7:129.
138. Thamm M, Balfanz S, Scheiner R, Baumann A, Blenau W. Characterization of the 5-HT1A receptor of the honeybee (*Apis mellifera*) and involvement of serotonin in phototactic behavior. *Cell Mol Life Sci* 2010; 67:2467-2479.
139. Troppmann B, Balfanz S, Baumann A, Blenau W. Inverse agonist and neutral antagonist actions of synthetic compounds at an insect 5-HT1 receptor. *Br J Pharmacol* 2010; 159:1450-1462.
140. Thamm M, Rolke D, Jordan N, Balfanz S, Schiffer C, Baumann A, et al. Function and distribution of 5-HT2 receptors in the honeybee (*Apis mellifera*). *PLoS One* 2013; 8:e82407.
141. Witz P, Amlaiky N, Plassat JL, Maroteaux L, Borrelli E, Hen R. Cloning and characterization of a *Drosophila* serotonin receptor that activates adenylyl cyclase. *Proc Natl Acad Sci U S A* 1990; 87:8940-8944.
142. Schlenstedt J, Balfanz S, Baumann A, Blenau W. Am5-HT7: molecular and pharmacological characterization of the first serotonin receptor of the honeybee (*Apis mellifera*). *J Neurochem* 2006; 98:1985-1998.
143. Vleugels R, Lenaerts C, Vanden Broeck J, Verlinden H. Signalling properties and pharmacology of a 5-HT7 -type serotonin receptor from *Tribolium castaneum*. *Insect Mol Biol* 2014; 23:230-243.
144. Pietrantonio PV, Jagge C, McDowell C. Cloning and expression analysis of a 5HT7-like serotonin receptor cDNA from mosquito *Aedes aegypti* female excretory and respiratory systems. *Insect Mol Biol* 2001; 10:357-369.
145. Röser C, Jordan N, Balfanz S, Baumann A, Walz B, Baumann O, et al. Molecular and pharmacological characterization of serotonin 5-HT2alpha and 5-HT7 receptors in the salivary glands of the blowfly *Calliphora vicina*. *PLoS ONE* 2012; 7:e49459.
146. Baumann O, Dames P, Kuhnelt D, Walz B. Distribution of serotonergic and dopaminergic nerve fibers in the salivary gland complex of the cockroach *Periplaneta americana*. *BMC Physiol* 2002; 2:9.
147. Baumann O, Kuhnelt D, Dames P, Walz B. Dopaminergic and serotonergic innervation of cockroach salivary glands: distribution and morphology of synapses and release sites. *J Exp Biol* 2004; 207:2565-2575.
148. Berridge MJ. The role of 5-hydroxytryptamine and cyclic AMP in the control of fluid secretion by isolated salivary glands. *J Exp Biol* 1970; 53:171-186.
149. Rietdorf K, Blenau W, Walz B. Protein secretion in cockroach salivary glands requires an increase in intracellular cAMP and Ca²⁺ concentrations. *J Insect Physiol* 2005; 51:1083-1091.
150. Molaei G, Lange AB. The association of serotonin with the alimentary canal of the African migratory locust, *Locusta migratoria*: distribution, physiology and pharmacological profile. *J Insect Physiol* 2003; 49:1073-1082.
151. Vanhoenacker P, Haegeman G, Leysen JE. 5-HT7 receptors: current knowledge and future prospects. *Trends Pharmacol Sci* 2000; 21:70-77.
152. Arvidsson LE, Hacksell U, Nilsson JL, Hjorth S, Carlsson A, Lindberg P, et al. 8-Hydroxy-2-(di-n-propylamino)tetralin, a new centrally acting 5-hydroxytryptamine receptor agonist. *J Med Chem* 1981; 24:921-923.
153. Krobert KA, Bach T, Syversveen T, Kvingedal AM, Levy FO. The cloned human 5-HT7 receptor splice variants: a comparative characterization of their pharmacology, function and distribution. *Naunyn Schmiedeberg's Arch Pharmacol* 2001; 363:620-632.
154. Thomas DR, Gittins SA, Collin LL, Middlemiss DN, Riley G, Hagan J, et al. Functional characterisation of the human cloned

- 5-HT₇ receptor (long form); antagonist profile of SB-258719. *Br J Pharmacol* 1998; 124:1300-1306.
155. Fitzgerald LW, Conklin DS, Krause CM, Marshall AP, Patterson JP, Tran DP, *et al.* High-affinity agonist binding correlates with efficacy (intrinsic activity) at the human serotonin 5-HT_{2A} and 5-HT_{2C} receptors: evidence favoring the ternary complex and two-state models of agonist action. *J Neurochem* 1999; 72:2127-2134.
156. Johnson M, Sonsalla PK, Letter AA, Hanson GR, Gibb JW. Role of the 5-HT₂ receptor in the methamphetamine-induced neurochemical alterations. *J Pharmacol Exp Ther* 1994; 270:97-103.
157. Kursar JD, Nelson DL, Wainscott DB, Baez M. Molecular cloning, functional expression, and mRNA tissue distribution of the human 5-hydroxytryptamine_{2B} receptor. *Mol Pharmacol* 1994; 46:227-234.
158. Newman-Tancredi A, Gavaudan S, Conte C, Chaput C, Touzard M, Verrielle L, *et al.* Agonist and antagonist actions of antipsychotic agents at 5-HT_{1A} receptors: a [³⁵S]GTPγS binding study. *Eur J Pharmacol* 1998; 355:245-256.
159. Newman-Tancredi A, Nicolas JP, Audinot V, Gavaudan S, Verrielle L, Touzard M, *et al.* Actions of α₂ adrenoceptor ligands at α_{2A} and 5-HT_{1A} receptors: the antagonist, atipamezole, and the agonist, dexmedetomidine, are highly selective for α_{2A} adrenoceptors. *Naunyn Schmiedeberg Arch Pharmacol* 1998; 358:197-206.
160. Lovell PJ, Bromidge SM, Dabbs S, Duckworth DM, Forbes IT, Jennings AJ, *et al.* A novel, potent, and selective 5-HT(7) antagonist: (R)-3-(2-(2-(4-methylpiperidin-1-yl)ethyl)pyrrolidine-1-sulfonyl)phenol (SB-269970). *J Med Chem* 2000; 43:342-345.
161. Thomas DR, Atkinson PJ, Ho M, Bromidge SM, Lovell PJ, Villani AJ, *et al.* [(3)H]-SB-269970--A selective antagonist radioligand for 5-HT(7) receptors. *Br J Pharmacol* 2000; 130:409-417.
162. Chemel BR, Roth BL, Armbruster B, Watts VJ, Nichols DE. WAY-100635 is a potent dopamine D₄ receptor agonist. *Psychopharmacology (Berl)* 2006; 188:244-251.
163. Foong JP, Bornstein JC. 5-HT antagonists NAN-190 and SB 269970 block α₂-adrenoceptors in the guinea pig. *Neuroreport* 2009; 20:325-330.
164. Yoshio R, Taniguchi T, Itoh H, Muramatsu I. Affinity of serotonin receptor antagonists and agonists to recombinant and native α₁-adrenoceptor subtypes. *Jpn J Pharmacol* 2001; 86:189-195.
165. Burris KD, Pacheco MA, Filtz TM, Kung MP, Kung HF, Molinoff PB. Lack of discrimination by agonists for D₂ and D₃ dopamine receptors. *Neuropsychopharmacology* 1995; 12:335-345.
166. Freedman SB, Patel S, Marwood R, Emms F, Seabrook GR, Knowles MR, *et al.* Expression and pharmacological characterization of the human D₃ dopamine receptor. *J Pharmacol Exp Ther* 1994; 268:417-426.
167. MacKenzie RG, VanLeeuwen D, Pugsley TA, Shih YH, Demattos S, Tang L, *et al.* Characterization of the human dopamine D₃ receptor expressed in transfected cell lines. *Eur J Pharmacol* 1994; 266:79-85.
168. Sunahara RK, Guan HC, O'Dowd BF, Seeman P, Laurier LG, Ng G, *et al.* Cloning of the gene for a human dopamine D₅ receptor with higher affinity for dopamine than D₁. *Nature* 1991; 350:614-619.
169. Lee DW, Pietrantonio PV. In vitro expression and pharmacology of the 5-HT₇-like receptor present in the mosquito *Aedes aegypti* tracheolar cells and hindgut-associated nerves. *Insect Mol Biol* 2003; 12:561-569.
170. Artigas F. 5-HT and antidepressants: new views from microdialysis studies. *Trends Pharmacol Sci* 1993; 14:262.
171. Celada P, Puig M, Amargos-Bosch M, Adell A, Artigas F. The therapeutic role of 5-HT_{1A} and 5-HT_{2A} receptors in depression. *J Psychiatry Neurosci* 2004; 29:252-265.
172. Celada P, Puig MV, Artigas F. Serotonin modulation of cortical neurons and networks. *Front Integr Neurosci* 2013; 7:25.
173. Marek GJ. Behavioral evidence for mu-opioid and 5-HT_{2A} receptor interactions. *Eur J Pharmacol* 2003; 474:77-83.
174. Quesseveur G, Nguyen HT, Gardier AM, Guiard BP. 5-HT₂ ligands in the treatment of anxiety and depression. *Expert Opin Investig Drugs* 2012; 21:1701-1725.
175. Vidal R, Diaz A, Pazos A, Castro E. Region-specific regulation of 5-HT_{1B} receptors in the rat brain by chronic venlafaxine treatment. *Psychopharmacology (Berl)* 2013; 229:177-185.
176. Mitsikostas DD, Tfelt-Hansen P. Targeting to 5-HT_{1F} receptor subtype for migraine treatment: lessons from the past, implications for the future. *Cent Nerv Syst Agents Med Chem* 2012; 12:241-249.
177. Briejer MR, Prins NH, Schuurkes JA. Effects of the enterokinetic prucalopride (R093877) on colonic motility in fasted dogs. *Neurogastroenterol Motil* 2001; 13:465-472.
178. Jandacek RJ, Anderson N, Liu M, Zheng S, Yang Q, Tso P. Effects of yo-yo diet, caloric restriction, and olestra on tissue distribution of hexachlorobenzene. *Am J Physiol Gastrointest Liver Physiol* 2005; 288:G292-G299.
179. Verlinden H, Vleugels R, Zels S, Dillen S, Lenaerts C, Crabbé K, *et al.* Neurohormone receptors as targets for the control of insect pest. *Adv Insect Physiol* 2014; 46:167-303.